Appl. No. 09/553,969
Amdt. dated November 25, 2008
Amendment under 37 CFR 1.116 Expedited Procedure
Examining Group 1611

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

- 1. (Currently Amended) An extrudable fragmented biocompatible resorbable single phase aqueous colloid which is substantially free from a free aqueous phase, said single phase aqueous colloid being present in an applicator having an extrusion orifice, wherein the single phase aqueous colloid has been fragmented by mechanical disruption, comprises a protein, has an equilibrium swell from 400% to 5000%, and has at least one characteristic selected from the group consisting of (a) a subunit size when fully hydrated in the range from 0.01 mm to 5 mm, (b) an equilibrium swell from 400% to 5000%; and (b) (e) an in vivo degradation time of less than one year.
 - 2-18. (Canceled)
- (Previously Presented) The single phase aqueous colloid of claim 1, having a subunit size when fully hydrated in the range from 0.01 mm to 5 mm.
 - 20. (Canceled)
- (Previously Presented) The single phase aqueous colloid of claim 1, having an in vivo degradation time of less than one year.
 - (Canceled)
 - (Canceled)
- 24. (Currently Amended) The single phase aqueous colloid of claim 1, wherein the single phase aqueous colloid has a subunit size when fully hydrated in the range from 0.01 mm to 5 mm and an in vivo degradation time of less than one year having all three characteristics.

Appl. No. 09/553,969
Amdt. dated November 25, 2008
Amendment under 37 CFR 1.116 Expedited Procedure
Examining Group 1611

- 25. (Previously Presented) The single phase aqueous colloid of claim 1, said single phase aqueous colloid being at least partially hydrated with an aqueous medium comprising an active agent.
- 26. (Previously Presented) The single phase aqueous colloid of claim 25, wherein the active agent is a clotting agent.
- (Previously Presented) The single phase aqueous colloid of claim 26, wherein the clotting agent is thrombin.
 - 28. (Canceled)
- (Currently Amended) The single phase aqueous colloid of claim <u>27</u>
 wherein the protein comprises gelatin.
- (Currently Amended) The single phase aqueous colloid of claim 27, wherein the <u>single phase aqueous colloid hydrogel</u> comprises a polysaccharide.
- (Previously Presented) The single phase aqueous colloid of claim 27, wherein the single phase aqueous colloid comprises a non-biological polymer.
- 32. (Currently Amended) The single phase aqueous colloid of claim 27, wherein the single phase aqueous colloid comprises two of the following components a) a protein, b) a polysaccharide or , and e) a non-biological polymer, or both.
- 33. (Currently Amended) The single phase aqueous colloid of claim 27, wherein the single phase aqueous colloid comprises a) a protein, b) a polysaccharide and e) a non-biological polymer.
- 34. (Currently Amended) An extrudable fragmented biocompatible resorbable single phase aqueous colloid present in an applicator having an extrusion orifice, wherein the single phase aqueous colloid has been fragmented by mechanical disruption, is not

Appl. No. 09/553,969
Amdt. dated November 25, 2008
Amendment under 37 CFR 1.116 Expedited Procedure
Examining Group 1611

hydrated above its capacity to absorb water, <u>has an equilibrium swell from 400% to 5000%</u>, and comprises gelatin, the single phase aqueous colloid having at least one characteristic selected from the group consisting of (a) a subunit size when fully hydrated in the range from 0.01 mm to 5 mm, (b) an equilibrium swell from 400% to 5000%, and (b) (e) an *in vivo* degradation time of less than one year.

- 35. (Currently Amended) An extrudable fragmented biocompatible resorbable single phase aqueous colloid which is substantially free from a free aqueous phase, said single phase aqueous colloid being present in an applicator having an extrusion orifice, wherein the single phase aqueous colloid has been fragmented by mechanical disruption, has an equilibrium swell from 400% to 5000%, and comprises a protein and a polysaccharide, the single phase aqueous colloid having at least one characteristic selected from the group consisting of (a) a subunit size when fully hydrated in the range from 0.01 mm to 5 mm, (b) an equilibrium swell from 400% to 5000%, and (b) (e) an in vivo degradation time of less than one year.
- 36. (Currently Amended) An extrudable fragmented biocompatible resorbable single phase aqueous colloid which is substantially free from a free aqueous phase, said single phase aqueous colloid being present in an applicator having an extrusion orifice, wherein the single phase aqueous colloid has been fragmented by mechanical disruption, has an equilibrium swell from 400% to 5000%, and comprises a protein and a non-biological polymer, the single phase aqueous colloid having at least one characteristic selected from the group consisting of (a) a subunit size when fully hydrated in the range from 0.01 mm to 5 mm, (b) an equilibrium swell from 400% to 5000%, and (b) (e) an in vivo degradation time of less than one year.